arXiv:1606.06860v1 [math.RA] 22 Jun 2016

Integral group rings with all central units trivial

Gurmeet K. Bakshi
Centre for Advanced Study in Mathematics,
Panjab University, Chandigarh 160014, India.
email: gkbakshi@pu.ac.in

Sugandha Maheshwary *†
Indian Institute of Science Education and Research, Mohali,
Sector 81, Mohali (Punjab)-140306, India.
email: sugandha@iisermohali.ac.in

Inder Bir S. Passi
Centre for Advanced Study in Mathematics,
Panjab University, Chandigarh-160014, India
&
Indian Institute of Science Education and Research, Mohali,
Sector 81, Mohali (Punjab)-140306, India.
email: ibspassi@yahoo.co.in

Abstract

The object of study in this paper is the finite groups whose integral group rings have only trivial central units. Prime-power groups and metacyclic groups with this property are characterized. Metacyclic groups are classified according to the central height of the unit groups of their integral group rings.

Keywords : integral group rings, unit group, trivial central units.

MSC2000 : 16U60; 16K20; 16S34; 20C05; 20F14

1 Introduction

Given a finite group G, let $\mathcal{U}(\mathbb{Z}[G])$ be the group of units of the integral group ring $\mathbb{Z}[G]$ and let $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G]))$ be its centre. The structure of $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G]))$ has been a

*Research supported by IISER, Mohali, India, is gratefully acknowledged
†Corresponding author

subject of intensive research (see e.g. [4, 10, 11, 14, 15, 16, 17, 18, 19, 25]). Clearly $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G]))$ contains the so-called *trivial central units* $\pm g$, $g \in \mathcal{Z}(G)$, the centre of G. Thus there arises the problem of characterizing the groups G having the property that all central units of the integral group ring $\mathbb{Z}[G]$ are trivial. For notational convenience, we refer to such groups as groups with the cut-property. *Abelian groups with the* cut*-property are precisely those having exponent* 1, 2, 3, 4 *or* 6 [12]. It has been shown by Ritter and Sehgal [25] that a finite group G has the cut-property if, and only if, for all $x \in G$ and for every natural number j, relatively prime to the order of G,

$$x^j \text{ is conjugate to either } x \text{ or } x^{-1}. \tag{1}$$

The above characterization is equivalent (see [25] & [13], p. 545) to saying that, if $\mathbb{Q}[G] \cong \bigoplus_i M_{n_i}(\mathbb{D}_i)$, is the Wedderburn decomposition of the rational group algebra $\mathbb{Q}[G]$, where $M_{n_i}(\mathbb{D}_i)$ denotes the algebra of $n_i \times n_i$ matrices over division ring $\mathbb{D}_i$, then

$$\text{the centre } \mathcal{Z}(\mathbb{D}_i) \text{ of each } \mathbb{D}_i \text{ is either } \mathbb{Q} \text{ or } \mathbb{Q}(\sqrt{-d}),\ d \in \mathbb{N}. \tag{2}$$

In particular, *the* cut*-property is quotient closed*, i.e., if G is a group with the cut-property, then so is every quotient of G. Our aim in this paper is to investigate further the cut-property for finite groups. The study of this property reveals, as expected, that it is a fairly restrictive property. After making some general observations on the groups with the cut-property, we examine this property in detail for nilpotent groups and metacyclic groups.

In Section 2, we prove (Theorem 1) that the order of a group with the cut-property must necessarily be divisible by 2 or 3. Consequently, since this property is quotient closed, a nilpotent group with the cut-property must be a (2, 3)-group. We note that the above-mentioned characterization (1) of groups with the cut-property can be considerably simplified in the case of 2-groups and 3-groups. We show that a 3-group G has the cut-property if, and only if, for all $x \in G$, x^2 is conjugate to x^{-1} (Theorem 2) and a 2-group G has the cut-property if, and only if, for each $x \in G$, x^3 is conjugate to either x or x^{-1} (Theorem 3). It is interesting to note that, for 3-groups, the cut-property is closed under direct sum, but it is not so for 2-groups.

In Section 3, as a contribution to understand the finite solvable groups with the cut-property, we examine metacyclic groups, and give a complete classification of

such groups with this property. It turns out that, up to isomorphism, only finitely many metacyclic groups have the cut-property, of which we give a complete list (Theorem 5).

It is known [1] that the central height of $V(\mathbb{Z}[G])$, the group of units of augmentation 1 in $\mathbb{Z}[G]$, is at most 2. We compute precisely the central height of $V(\mathbb{Z}[G])$, when G is a metacyclic group (Theorem 6).

2 Finite groups with $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G])) = \pm\mathcal{Z}(G)$

We begin by observing that the cut-property has strong bearing on the centre of the group. If G has the cut-property, then so does its centre $\mathcal{Z}(G)$. For if $\mathcal{Z} = \mathcal{Z}(G)$, then $\mathcal{U}(\mathbb{Z}[\mathcal{Z}]) \subseteq \mathcal{Z}(\mathcal{U}(\mathbb{Z}[G]))$. In case G has the cut-property, $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G])) = \pm\mathcal{Z}(G)$, and consequently, in that case, $\mathcal{U}(\mathbb{Z}[\mathcal{Z}]) = \pm\mathcal{Z}$.

For a finite group G, let $\pi(G)$ denote the set of primes which divide the order of G.

Theorem 1 *If G has the* cut*-property, then either 2 or $3 \in \pi(G)$. Furthermore, if G is nilpotent, then $\pi(G) \subseteq \{2, 3\}$.*

Proof. Let G be a group with the cut-property. If $2 \notin \pi(G)$, then, by the odd-order theorem [9], G is solvable and hence $G \neq G'$, the derived subgroup of G. Let p be a prime divisor of $[G : G']$, the index of G' in G. Since the cut-property is quotient-closed, G/G' has the cut-property. Thus it follows that G/G' is a 3-group, and hence $3 \in \pi(G)$. Further, if G has the cut-property, then so does $G/\mathcal{Z}(G)$. Thus, if G is nilpotent group, then each quotient in the upper central series of G, being abelian with the cut-property, is of exponent $2, 3, 4$ or 6 and hence $\pi(G) \subseteq \{2, 3\}$. $\square$

We next investigate the 2-groups and 3-groups with the cut-property.

Theorem 2 *A finite 3-group has the* cut*-property if, and only if, x^2 is conjugate to x^{-1}, for all $x \in G$. Furthermore, if a 3-group G has the* cut*-property, then each quotient in the lower, as well as the upper, central series of G is of exponent 3.*

Proof. In view of the characterization (1) and the fact that 2 is a primitive root modulo 3^n, for all $n \geq 1$, a 3-group G has the cut-property if, and only if, x^2 is conjugate to x or x^{-1}, for all $x \in G$.

Let G be a 3-group and let

$$G = \gamma_1(G) \supset \gamma_2(G) \supset ... \supset \gamma_n(G) = \langle 1 \rangle$$

be its lower central series. We claim that x^2 is not conjugate to x for any $1 \neq x \in G$. For, given $1 \neq x \in G$, there exists i $(1 \leq i < n)$ such that $x \in \gamma_i(G) \setminus \gamma_{i+1}(G)$. If x^2 were conjugate to x, then $x^2 = g^{-1}xg$ for some $g \in G$, and therefore $x = x^{-1}g^{-1}xg \in \gamma_{i+1}(G)$, which contradicts the assumption. This proves the first statement.

Thus, if G is a 3-group with the cut-property, then for every $1 \neq x \in G$ and $1 \leq j < n$, there exits $y \in G$ such that $x^2 = y^{-1}x^{-1}y$ and therefore, $x^3 \in \gamma_{j+1}(G)$.

Next, let

$$\langle 1 \rangle = \mathcal{Z}_0(G) \subset \mathcal{Z}_1(G) \subset ... \subset \mathcal{Z}_n(G) = G$$

be the upper central series of G. Then, $\mathcal{Z}_1(G) = \mathcal{Z}(G)$ is of exponent 3 and as the cut-property is quotient closed, it follows inductively that, for every i, $\mathcal{Z}_{i+1}(G)/\mathcal{Z}_i(G)$ is of exponent 3. □

Corollary 1 *The* cut*-property is direct sum closed for* 3*-groups.*

Proof. Let H and K be 3-groups with the cut-property and let $G = H \times K$ be the direct sum of H and K. Further, let $g = (h, k) \in G$, $h \in H$, $k \in K$. Then, $g^2 = (h^2, k^2)$ is conjugate to $(h^{-1}, k^{-1}) = g^{-1}$ and hence, by Theorem 2, G has the cut-property. □

In view of Eq. (1) and the fact that $\mathcal{U}(\mathbb{Z}/2^n\mathbb{Z}) = \langle \pm 1 \rangle \oplus \langle 3 \rangle$, we have the following analogous result for 2-groups.

Theorem 3 *A finite* 2*-group has the* cut*-property if, and only if, x^3 is conjugate to x or x^{-1}, for all $x \in G$. Furthermore, if a* 2*-group G has the* cut*-property, then each quotient in its lower central, as well as the upper central, series is of exponent* 2 *or* 4*.*

Remark 1 *Direct sum of* 2*-groups with the* cut*-property does not necessarily have the* cut*-property. For example, let*

$$H = \langle a, b \mid a^8 = b^2 = 1, b^{-1}ab = a^3 \rangle \text{ and } K = \langle x \mid x^4 = 1 \rangle.$$

It is easy to check that both H and K have the cut*-property. However, $g = (a, x) \in H \times K$, is such that $g^3 = (a^3, x^3)$ is neither conjugate to g nor to g^{-1}.*

An interesting class of p-groups with the cut-property is provided by Camina groups. Recall that a group G is called a *Camina group*, if $G' \neq G$ and, for every

$g \notin G'$, the coset gG' is a conjugacy class. For a normal subgroup N of G, the pair (G, N) is said to be a *Camina pair*, if each element $g \notin N$ is conjugate to all the elements in the coset gN, so that G is a Camina group, if (G, G') is a Camina pair. Camina p-groups have nilpotency class at most three, and the factors of their lower central series are of exponent p [8, 21].

Theorem 4 *Non-abelian Camina p-groups have the* cut*-property for $p = 2,\ 3$.*

Proof. Let G be a Camina 3-group of class 2. Then G/G' and G' are both of exponent 3. It thus immediately follows that x^2 is conjugate to x^{-1} for every $x \in G$, and consequently, by Theorem 2, G has the cut-property.

In case G is a Camina 2-group of class 2, then G has exponent 4, and hence $x^3 = x^{-1}$ for every $x \in G$. Thus, in view of Theorem 3, G has the cut-property.

Next, let G be a p-group of nilpotency class 3, $p = 2,\ 3$. Then $G/\gamma_3(G)$ is a Camina p-group of class 2, and so it has the cut-property. Furthermore, $(G, \gamma_3(G))$ forms a Camina pair ([21], Theorem 5.2), which ensures, by Theorems 2 & 3, that G has the cut-property. □

3 Finite metacyclic groups

3.1 Metacyclic groups with the cut-property

In this section, we determine the metacyclic groups with the cut-property.
We begin by recalling some results from [22] .

Let G be a finite group and K be a normal subgroup of subgroup H of G. Define $\hat{H} := \frac{1}{|H|}\sum_{h\in H} h$ and

$$\varepsilon(H,K) := \begin{cases} \hat{H}, & H = K; \\ \prod(\hat{K} - \hat{L}), & \text{otherwise,} \end{cases}$$

where $|H|$ denotes the order of H and L runs over the normal subgroups of H, which are minimal over the normal subgroups of H containing K properly.

A *strong Shoda pair* of G is a pair (H, K) of subgroups of G with the property that

(i) K is normal in H and H is normal in $N_G(K)$, the normalizer of K in G;

(ii) H/K is cyclic and a maximal abelian subgroup of $N_G(K)/K$;

(iii) the distinct G-conjugates of $\varepsilon(H, K)$ are mutually orthogonal.

It is known that if (H, K) is a strong Shoda pair of G, then $e(G, H, K)$, the sum of distinct G-conjugates of $\varepsilon(H, K)$, is a primitive central idempotent of $\mathbb{Q}[G]$ ([22], Proposition 3.3).

Furthermore, a strong Shoda pair (H, K) of G with H normal in G has an easier description in view of the following:

Proposition 1 ([22], Proposition 3.6) *Let (H, K) be a pair of subgroups of a finite group G satisfying the following conditions:*

(i) *K is normal in H and H is normal in G,*

(ii) *H/K is cyclic and a maximal abelian subgroup of $N_G(K)/K$.*

Then (H, K) is a strong Shoda pair and hence $e(G, H, K)$ is a primitive central idempotent of $\mathbb{Q}[G]$.

A big advantage of determining the primitive central idempotent of $\mathbb{Q}[G]$, using strong Shoda pair (H, K) of G is that one can describe the structure of the simple component $\mathbb{Q}[G]e(G, H, K)$ of $\mathbb{Q}[G]$, corresponding to the primitive central idempotent $e(G, H, K)$, in view of the next proposition.

Note that $R *_\tau^\sigma G$ denotes the crossed product of group G over the coefficient ring R with action σ and twisting τ [23]. Recall that a classical crossed product is a crossed product $K *_\tau^\sigma G$, where K/F is a finite Galois extension, $G = \mathrm{Gal}(K/F)$ is the Galois group of K/F and σ is the natural action of G on K. The classical crossed product is denoted by $(K/F, \tau)$ [24].

Proposition 2 ([22], Proposition 3.4) *Let (H, K) be a strong Shoda pair of a finite group G with $k = [H : K]$, $N = N_G(K)$, $n = [G : N]$, x a generator of H/K and $\phi : N/H \mapsto N/K$ a left inverse of the projection $N/K \mapsto N/H$. Then,*

$$\mathbb{Q}[G]e(G, H, K) \cong M_n(\mathbb{Q}(\zeta_k) *_\tau^\sigma N/H),$$

where ζ_k denotes the k^{th} root of unity, the action σ and the twisting τ are given by $\zeta_k^{\sigma(a)} = \zeta_k^i$, if $x^{\phi(a)} = x^i$; $\tau(a, b) = \zeta_k^j$, if $\phi(ab)^{-1}\phi(a)\phi(b) = x^j$, for $a, b \in N/H$ and integers i and j.

Remark 2 *The action σ in the above proposition is faithful and therefore $\mathbb{Q}(\zeta_k) *_\tau^\sigma N/H$ can be described as a classical crossed product $(\mathbb{Q}(\zeta_k)/F, \tau)$, where*

F is the centre of the algebra and σ is determined by the action of N/H on H/K. In this way, the Galois group $\mathrm{Gal}(\mathbb{Q}(\zeta_k)/F)$ *can be identified with N/H and with this identification, $F = \mathbb{Q}(\zeta_k)^{N/H}$, the fixed field of N/H.*

We now state the main theorem of this section, which classifies all metacyclic groups with the cut-property.

Theorem 5 *Let G be a finite non-abelian metacyclic group given by*

$$G = \langle a,\ b \mid a^n = 1,\ b^t = a^\ell,\ b^{-1}ab = a^r \rangle, \tag{3}$$

where n, t, r, ℓ are natural numbers such that $r^t \equiv 1 \pmod{n}$, $\ell r \equiv \ell \pmod{n}$ and ℓ divides n. Then, G has the cut*-property if, and only if, G is isomorphic to one of the following:*

$\langle a,\ b \mid a^n = 1,\ b^t = 1,\ b^{-1}ab = a^{n-1} \rangle,\quad t = 2, 4, 6,\quad n = 3, 4, 6;$
$\langle a,\ b \mid a^4 = 1,\ b^t = a^2,\ b^{-1}ab = a^3 \rangle,\ t = 2, 4, 6;$
$\langle a,\ b \mid a^6 = 1,\ b^2 = a^3,\ b^{-1}ab = a^5 \rangle;$
$\langle a,\ b \mid a^n = 1,\ b^{\varphi(n)} = 1,\ b^{-1}ab = a^{\lambda_n} \rangle,\ n = 5, 7, 9, 10, 14, 18;$
$\langle a,\ b \mid a^n = 1,\ b^{\frac{\varphi(n)}{j}} = 1,\ b^{-1}ab = a^{\lambda_n^2} \rangle,\quad j = 1, 2,\quad n = 7, 9, 14, 18;$
$\langle a,\ b \mid a^8 = 1,\ b^t = 1,\ b^{-1}ab = a^r \rangle,\ t = 2, 4,\ r = 3, 5;$
$\langle a,\ b \mid a^{12} = 1,\ b^t = 1,\ b^{-1}ab = a^5 \rangle,\ t = 2, 4;$
$\langle a,\ b \mid a^{12} = 1,\ b^t = a^\ell,\ b^{-1}ab = a^7 \rangle,\ t = 2, 6,\ \ell = t, 12;$
$\langle a,\ b \mid a^{15} = 1,\ b^4 = 1,\ b^{-1}ab = a^2 \rangle;$
$\langle a,\ b \mid a^{16} = 1,\ b^4 = 1,\ b^{-1}ab = a^r \rangle,\ r = 3, 5;$
$\langle a,\ b \mid a^{20} = 1,\ b^4 = 1,\ b^{-1}ab = a^r \rangle,\ r = 3, 13;$
$\langle a,\ b \mid a^{20} = 1,\ b^4 = a^{10},\ b^{-1}ab = a^3 \rangle;$
$\langle a,\ b \mid a^{21} = 1,\ b^6 = 1,\ b^{-1}ab = a^r \rangle,\ r = 2, 10;$
$\langle a,\ b \mid a^{28} = 1,\ b^6 = a^\ell,\ b^{-1}ab = a^{11} \rangle,\ \ell = 14, 28;$
$\langle a,\ b \mid a^{30} = 1,\ b^4 = 1,\ b^{-1}ab = a^{17} \rangle;$
$\langle a,\ b \mid a^{36} = 1,\ b^6 = a^\ell,\ b^{-1}ab = a^7 \rangle,\ \ell = 6, 36;$
$\langle a,\ b \mid a^{42} = 1,\ b^6 = 1,\ b^{-1}ab = a^r \rangle,\ r = 11, 19;$

where λ_n is a generator of $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ and φ denotes the Euler's phi function.

Proof. Let G be as in (3). Suppose G has the cut-property. Since the cut-property is quotient closed, we have that the cyclic group $G/\langle a \rangle$ also has the cut-property and hence,

$$t = 2, 3, 4 \text{ or } 6. \tag{4}$$

Denote by $o_n(r)$, the order of r modulo n. If $o_n(r) = d$, then one can check that $(\langle a,\, b^d\rangle,\ \langle a^\alpha b^d\rangle)$ is a strong Shoda pair of G, for some $0 \leq \alpha < n$. For instance, the following may be a choice of α for various cases:

- If $t = d$, then $\alpha = n - \ell$.
- If $t \neq d$ but $\ell = n$, then $\alpha = n$.
- If $t \neq d$, $\ell \neq n$ and $\mathrm{g.c.d.}(\frac{t}{d}, n)$ divides ℓ, where $\mathrm{g.c.d.}(\frac{t}{d}, n)$ denotes the greatest common divisor of $\frac{t}{d}$ and n, then α is a solution of $\frac{t}{d}x \equiv -\ell \pmod{n}$.
- For all the other cases, $\alpha = \begin{cases} \frac{m-\ell}{2}, & \text{if } \frac{t}{d} = 2; \\ \frac{m-\ell}{3} \text{ or } \frac{2m-\ell}{3}, & \text{if } \frac{t}{d} = 3, \end{cases}$ where m denotes the greatest divisor of n, which is co-prime to $\frac{t}{d}$.

In view of (2), the centre of the simple component of $\mathbb{Q}[G]$ corresponding to this strong Shoda pair, computed using Proposition 2, must be either $\mathbb{Q}$ or imaginary quadratic. Therefore, we must have

$$\frac{\varphi(n)}{d} \leq 2.$$

Consequently, $[\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) : \langle r\rangle] \leq 2$. We consider the following possible cases:

Case(I) $\underline{[\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) : \langle r\rangle] = 1}$

Since $\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) = \langle r\rangle$, $o_n(r)$ divides t and $t = 2, 3, 4$ or 6 (by (4)), we have in this case $o_n(r) = \varphi(n) = 2, 3, 4$ or 6.

Subcase (i) $\underline{\varphi(n) = 2}$. Thus, $n = 3, 4, 6$ and $r = n - 1$. In this case, for any n, the admissible values of t are $2, 4, 6$. Therefore, for suitable choices of ℓ, $G = \langle a,\ b \mid a^n = 1,\ b^t = a^\ell,\ b^{-1}ab = a^{n-1}\rangle$, where $n = 3, 4$ or 6 and $t = 2, 4$ or 6.

Subcase (ii) $\underline{\varphi(n) \neq 2}$. Observe that in this case, $t = o_n(r) = \varphi(n) = 4$ or 6. As $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ is cyclic, $n = 5, 7, 9, 10, 14$ or 18. Hence, for suitable choices of ℓ, $G \cong \langle a,\ b \mid a^n = 1,\ b^t = a^\ell,\ b^{-1}ab = a^{\lambda_n}\rangle$, where $n = 5, 7, 9, 10, 14$ or 18 and $t = \varphi(n)$.

Note that if n is even and ℓ is odd, then $t \neq 4$, 6. For otherwise, $(G, \langle a^2\rangle)$ is a strong Shoda pair of G and centre of the simple component corresponding to this pair is $\mathbb{Q}(\zeta_{2t}) = \mathbb{Q}(\zeta_8)$ or $\mathbb{Q}(\zeta_{12})$. Therefore, we get that any group in this case, which has the cut-property, must be one of the following:

$\langle a,\ b \mid a^n = 1,\ b^t = 1,\ b^{-1}ab = a^{n-1}\rangle,\quad t = 2, 4, 6,\quad n = 3, 4, 6;$
$\langle a,\ b \mid a^4 = 1,\ b^t = a^2,\ b^{-1}ab = a^{n-1}\rangle,\ t = 2, 4, 6;$
$\langle a,\ b \mid a^6 = 1,\ b^2 = a^3,\ b^{-1}ab = a^{n-1}\rangle;$
$\langle a,\ b \mid a^n = 1,\ b^{\varphi(n)} = 1,\ b^{-1}ab = a^{\lambda_n}\rangle,\ n = 5, 7, 9, 10, 14, 18.$

Case(II) $[\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) : \langle r\rangle] = 2$ **and** $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ **is cyclic**

Clearly, $o_n(r) = \frac{\varphi(n)}{2}$. As $o_n(r)$ divides t, $\varphi(n) = 2, 4, 6, 8$ or 12. Since G is non abelian, $\varphi(n) \neq 2$. Also, $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ is cyclic yields following subcases:

Subcase(i) $\varphi(n) = 4$, $n = 5, 10$. In this case $t = 2,\ 4$ or 6; and $r = n - 1$. Notice that the centre of the simple component corresponding to the strong Shoda pair $(\langle a, b^2\rangle, \langle b^2\rangle)$ is $\mathbb{Q}(\zeta_n + \zeta_n^{-1})$, which is a real quadratic extension of $\mathbb{Q}$. Hence, this case yields no group with the cut-property.

Subcase(ii) $\varphi(n) = 6$, $n = 7, 9, 14, 18$. In this case, $t = 3$ or 6, as $o_n(r)$ divides t. As noted in Case(I), if $n = 14$ and $t = 6$, then $\ell \neq 7$ and hence, we must have $b^t = 1$ i.e. $G \cong \langle a,\ b \mid a^n = 1,\ b^{\varphi(n)} = 1,\ b^{-1}ab = a^{\lambda_n^2}\rangle$ or $G \cong \langle a,\ b \mid a^n = 1,\ b^{\frac{\varphi(n)}{2}} = a^\ell,\ b^{-1}ab = a^{\lambda_n^2}\rangle$, $\ell = 0 \pmod 7$, if $n = 7, 14$. Similar arguments yield that $\ell \equiv 0 \pmod 6$, if $n = 18$ and $t = 6$. Otherwise, $\ell \equiv 0 \pmod 3$, for $n = 9, 18$. Further, in view of [6], the groups listed above are isomorphic to one of the following:

$$\langle a,\ b \mid a^n = 1,\ b^{\frac{\varphi(n)}{j}} = 1,\ b^{-1}ab = a^{\lambda_n^2}\rangle,\quad j = 1, 2,\quad n = 7, 9, 14, 18. \quad (5)$$

Subcase(iii) $\varphi(n) = 12$, $n = 13, 26$. Here, $(\langle a\rangle, \langle 1\rangle)$ is a strong Shoda pair and the centre of the corresponding simple component is contained in $\mathbb{Q}(\zeta_n + \zeta_n^{-1})$, which is a real extension of $\mathbb{Q}$. Hence, this case also yields no group with the cut-property.

Therefore, if $[\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) : \langle r\rangle] = 2$ and $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ is cyclic group generated by λ_n, then G is isomorphic to one of the groups in (5).

Case(III) $[\mathcal{U}(\mathbb{Z}/n\mathbb{Z}) : \langle r\rangle] = 2$ **and** $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ **is non-cyclic**

For each group in this case, we enlist below the strong Shoda pair of G, denoted by *SSP*, which yields the simple component of $\mathbb{Q}[G]$, whose centre is different from $\mathbb{Q}$ or imaginary quadratic and hence restricts G from having the cut-property. For each n, we list the choice of parameters t, r and ℓ, which define the group G, as given in (3). As $\mathcal{U}(\mathbb{Z}/n\mathbb{Z})$ is non-cyclic, $o_n(r) = \frac{\varphi(n)}{2}$ and $o_n(r)$ divides t, we have the following subcases:

Subcase(i) $\varphi(n) = 4,\ \ n = 8, 12,\ \ t = 2, 4$ or $6.$

$\underline{n = 8}$

t	r	ℓ	*SSP*
2	7	4,8	$(\langle a\rangle,\ \langle 1\rangle)$
4	3	4	$(\langle a,\ b^2\rangle,\ \langle a^2b^2\rangle)$
4	5	2	$(\langle a,\ b^2\rangle,\ \langle a^3b^2\rangle)$
4	7	8	$(\langle a,\ b^2\rangle,\ \langle b^2\rangle)$
6	3,7	8	$(\langle a,\ b^2\rangle,\ \langle b^2\rangle)$
6	5	8	$(\langle a,\ b\rangle,\ \langle a^4,\ b^3\rangle)$
6	5	4	$(\langle a,\ b^2\rangle,\ \langle a^4b^2\rangle)$

$\underline{n = 12}$

t	r	ℓ	*SSP*
2	5	3	$(\langle a,\ b\rangle,\ \langle a^4\rangle)$
2	11	6,12	$(\langle a\rangle,\ \langle 1\rangle)$
4	5	3	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
4	5	6	$(\langle a,\ b\rangle,\ \langle ab^2\rangle)$
4	7	2,12	$(\langle a,\ b\rangle,\ \langle a^3\rangle)$
4	11	12	$(\langle a,\ b^2\rangle,\ \langle b^2\rangle)$
4	11	6	$(\langle a,\ b^2\rangle,\ \langle a^9b^2\rangle)$
6	5	12	$(\langle a,\ b\rangle,\ \langle a^4,\ b^3\rangle)$
6	5	3	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
6	7	2,4	$(\langle a,\ b\rangle,\ \langle a^3\rangle)$
6	11	12	$(\langle a,\ b^2\rangle,\ \langle a^4b^2\rangle)$
6	11	6	$(\langle a,\ b^2\rangle,\ \langle a^{10}b^2\rangle)$

Subcase(ii) $\varphi(n) = 8,\ \ n = 15, 16, 20, 30,\ \ t = 4.$

$\underline{n = 15}$

t	r	ℓ	*SSP*
4	7	15	$(\langle a,\ b\rangle,\ \langle a^3\rangle)$

$\underline{n = 20}$

t	r	ℓ	*SSP*
4	13	5	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
4	13	10	$(\langle a,\ b\rangle,\ \langle ab^2,\ a^4\rangle)$

$\underline{n = 30}$

t	r	ℓ	*SSP*
4	7	5,30	$(\langle a,\ b\rangle,\ \langle a^3\rangle)$
4	17	15	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$

Subcase(iii) $\varphi(n) = 12,\ \ n = 21, 28, 36, 42,\ \ t = 6.$

$\underline{n = 21}$

t	r	ℓ	*SSP*
6	5	21	$(\langle a\rangle,\ \langle 1\rangle)$
6	10	7	$(\langle a,\ b\rangle,\ \langle a^{15}\rangle)$

$\underline{n = 28}$

t	r	ℓ	*SSP*
6	3	14,28	$(\langle a\rangle,\ \langle 1\rangle)$
6	5	28	$(\langle a,\ b\rangle,\ \langle a^4,\ b^3\rangle)$
6	5	7	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$

$\underline{n = 36}$

t	r	ℓ	*SSP*
6	5	36	$(\langle a,\ b\rangle,\ \langle a^4,\ b^3\rangle)$
6	5	9	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
6	11	36	$(\langle a,\ b^2\rangle,\ \langle a^4b^2\rangle)$
6	11	18	$(\langle a,\ b^2\rangle,\ \langle a^{10}b^2\rangle)$

$\underline{n = 42}$

t	r	ℓ	*SSP*
6	5	42	$(\langle a\rangle,\ \langle 1\rangle)$
6	5, 11	21	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
6	19	7,21	$(\langle a,\ b\rangle,\ \langle a^2\rangle)$
6	19	14	$(\langle a,\ b\rangle,\ \langle a^3\rangle)$

Clearly, the groups in subcases (i)-(iii) (with different set of parameters) which are isomorphic to one of the groups tabulated above, fail to have the cut-property. The remaining ones are isomorphic to one of the following:

$\langle a,\ b \mid a^8 = 1,\ b^t = 1,\ b^{-1}ab = a^r\rangle,\ t = 2, 4,\ r = 3, 5.$
$\langle a,\ b \mid a^{12} = 1,\ b^t = 1,\ b^{-1}ab = a^5\rangle,\ t = 2, 4.$
$\langle a,\ b \mid a^{12} = 1,\ b^t = a^\ell,\ b^{-1}ab = a^7\rangle,\ t = 2, 6,\ \ell = t, 12.$
$\langle a,\ b \mid a^{15} = 1,\ b^4 = 1,\ b^{-1}ab = a^2\rangle.$
$\langle a,\ b \mid a^{16} = 1,\ b^4 = 1,\ b^{-1}ab = a^r\rangle,\ r = 3, 5.$
$\langle a,\ b \mid a^{20} = 1,\ b^4 = 1,\ b^{-1}ab = a^r\rangle,\ r = 3, 13.$
$\langle a,\ b \mid a^{20} = 1,\ b^4 = a^{10},\ b^{-1}ab = a^3\rangle.$
$\langle a,\ b \mid a^{21} = 1,\ b^6 = 1,\ b^{-1}ab = a^r\rangle,\ r = 2, 10.$
$\langle a,\ b \mid a^{28} = 1,\ b^6 = a^\ell,\ b^{-1}ab = a^{11}\rangle,\ \ell = 14, 28.$
$\langle a,\ b \mid a^{30} = 1,\ b^4 = 1,\ b^{-1}ab = a^{17}\rangle.$
$\langle a,\ b \mid a^{36} = 1,\ b^6 = a^\ell,\ b^{-1}ab = a^7\rangle,\ \ell = 6, 36.$
$\langle a,\ b \mid a^{42} = 1,\ b^6 = 1,\ b^{-1}ab = a^r\rangle,\ r = 11, 19.$

For isomorphism of groups, we have again used the results of [6].

It only remains to check that each group G listed in the theorem has the cut-property. Since metacyclic groups are normally monomial [5], and hence strongly monomial [3], this can be done using Theorem 3.1 of [15] (see also [4], Theorem 2) on the rank of the free abelian component of $\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G]))$ and Theorem 1 of [3] on the computation of strong Shoda pairs of G. $\square$

Remark 3 *In view of* ([13], p. 545)*, an alternative approach to determine metacyclic groups with the* cut*-property would be to compute the character fields of their complex irreducible representations* [7]*.*

Remark 4 *Recall that a group G is said to have the normalizer property, if $G.\mathcal{Z}(\mathcal{U}(\mathbb{Z}[G])) = N_{\mathcal{U}}(G)$, the normalizer of G in $\mathcal{U}(\mathbb{Z}[G])$. It is known* ([20], Lemma 2.1) *that the* cut*-property implies the normalizer property. It is interesting to note that although it is easy to see that all metacyclic groups have the normalizer property, only a few of them have the cut property.*

3.2 The upper central series

It is well known ([1], Theorem 2.6) that, for a finite group G, the central height of $V(\mathbb{Z}[G])$, the group of units of augmentation 1 in $\mathbb{Z}[G]$, is at most 2. In this section, we give a classification of finite metacyclic groups G for the permissible values of the central height of $V(\mathbb{Z}[G])$.

Theorem 6 *Let G be a finite metacyclic group and let $V(\mathbb{Z}[G])$ be the group of units of augmentation 1 in $\mathbb{Z}[G]$.*

(I) *The central height of $V(\mathbb{Z}[G])$ equals 0 if, and only if, G is isomorphic to one of following:*

$$\langle a,\ b \mid a^3 = 1,\ b^2 = 1,\ b^{-1}ab = a^2\rangle$$
$$\langle a,\ b \mid a^5 = 1,\ b^4 = 1,\ b^{-1}ab = a^2\rangle$$
$$\langle a,\ b \mid a^7 = 1,\ b^6 = 1,\ b^{-1}ab = a^3\rangle$$
$$\langle a,\ b \mid a^9 = 1,\ b^6 = 1,\ b^{-1}ab = a^2\rangle$$
$$\langle a,\ b \mid a^7 = 1,\ b^3 = 1,\ b^{-1}ab = a^2\rangle$$
$$\langle a,\ b \mid a^{15} = 1,\ b^4 = 1,\ b^{-1}ab = a^2\rangle$$
$$\langle a,\ b \mid a^{21} = 1,\ b^6 = 1,\ b^{-1}ab = a^2\rangle.$$

(II) *The central height of $V(\mathbb{Z}[G])$ equals 2 if, and only if, G is isomorphic to*

$$\langle a,\ b \mid a^n = 1,\ b^2 = a^{\frac{n}{2}},\ b^{-1}ab = a^{-1}\rangle,$$

where n is a positive integer divisible by 4.

(III) *For all other groups G, the central height of $V(\mathbb{Z}[G])$ equals 1.*

Proof. Let G be as in (3).

(I) Clearly, if the central height of $V(\mathbb{Z}[G])$ equals 0, i.e., $V(\mathbb{Z}[G]) = \langle 1 \rangle$, then G must be one of the groups listed in Theorem 5 with $\mathcal{Z}(G) = \langle 1 \rangle$. Observe that $1 \neq a^i b^j \in \mathcal{Z}(G)$ if, and only if, for some i, j, not both 0, $ri \equiv i (\text{mod } n)$ and $r^j \equiv 1 (\text{mod } n)$, where $0 \leq i < n$ and $0 \leq j < t$. This occurs if, and only if, either $\text{g. c. d.}(r-1, n) \neq 1$ or $b^{o_n(r)} \neq 1$. These observations yield (I).

(II) In view of ([2], Theorem 3.7), the central height of $V(\mathbb{Z}[G])$ is 2 if, and only if, G is a Q^* group, i.e., there exists an abelian subgroup H of G of index 2 and an element x of order 4 such that $G = \langle H, x \rangle$ and the following holds:

(i) $x^{-1}hx = h^{-1}$ for each $h \in H$;
(ii) $x^2 = y^2$ for some $y \in H$;
(iii) H is not an elementary Abelian 2-group.

We thus examine which metacyclic groups are Q^* groups. As the index of H in G is 2, we have that $a^2 \in H$, and therefore $H \cap \langle a \rangle = \langle a \rangle$ or $\langle a^2 \rangle$. We take these two cases seperately.

$\underline{H \cap \langle a \rangle = \langle a \rangle.}$
The index of H in G being 2, we must have that $H = \langle a, b^2 \rangle$. Furthermore, as H is abelian, b^2 commutes with a and hence $b^2 \in \mathcal{Z}(G)$. This gives $b^4 = 1$ since $x^{-1}b^2x = b^{-2}$ (by (i)). As $x \in Hb$, i.e., $x = h'b$, for some $h' \in H$, it follows from (i) that $b^{-1}hb = h^{-1}$, for every $h \in H$ and hence $x^2 = h'bh'b = b^2$. This implies that the order of b must be 4. Also by (ii), there exists an element $y \in H$ of order 4 with $x^2 = y^2$. However, H has an element of order 4 if, and only if, 4 divides n, and in that case, the only elements in H of order 4 are $(a^{\frac{n}{4}})^i b^{2j}$, with $i = 1, 3$ and $j = 0, 1$. Further, notice that the square of all of these four elements is $a^{\frac{n}{2}}$. Consequently, we must have $b^2 = a^{\frac{n}{2}}$ and $G \cong \langle a, b \mid a^n = 1,\ b^2 = a^{\frac{n}{2}},\ b^{-1}ab = a^{-1} \rangle$, where n is divisible by 4.

$\underline{H \cap \langle a \rangle = \langle a^2 \rangle.}$
In this case, we have $H = \langle a^2, b \rangle$ or $\langle a^2, ab \rangle$, depending upon whether b belongs to H or not. Suppose $H = \langle a^2, b \rangle$. Since H is abelian, we have that a^2 commutes with b, so that $a^2 \in \mathcal{Z}(G)$. It thus follows by (i) that $a^4 = 1$. However, $\langle a \rangle$ is a normal subgroup of non abelian group G, which yields that the order of a cannot be 2. Therefore, the order of a is 4 and bab^{-1} must be a^{-1}. Further as $x \in Ha$, $x^{-1}bx = b^{-1}$ implies that $a^{-1}ba = b^{-1}$. Consequently, $a^2 = b^2$ and $G \cong \langle a, b \mid a^4 = 1,\ b^2 = a^2,\ b^{-1}ab = a^{-1} \rangle$.

Similarly, if $H = \langle a^2, ab \rangle$, then $G \cong \langle a, b \mid a^4 = 1,\ b^2 = a^2,\ b^{-1}ab = a^{-1} \rangle$. $\square$

References

[1] S. R. Arora, A. W. Hales, and I. B. S. Passi, *Jordan decomposition and hypercentral units in integral group rings*, Comm. Algebra **21** (1993), no. 1, 25–35.

[2] S. R. Arora and I. B. S. Passi, *Central height of the unit group of an integral group ring*, Comm. Algebra **21** (1993), no. 10, 3673–3683.

[3] G.K. Bakshi and S. Maheshwary, *The rational group algebra of a normally monomial group*, J. Pure Appl. Algebra **218** (2014), no. 9, 1583–1593.

[4] ———, *On the index of a free abelian subgroup in the group of central units of an integral group ring*, J. Algebra **434** (2015), 72–89.

[5] B. G. Basmaji, *Monomial representations and metabelian groups*, Nagoya Math. J. **35** (1969), 99–107.

[6] ———, *On the ismorphisms of two metacyclic groups*, Proc. Amer. Math. Soc. **22** (1969), 175–182.

[7] ———, *Complex representations of metacyclic groups*, Amer. Math. Monthly **86** (1979), no. 1, 47–48.

[8] R. Dark and C. M. Scoppola, *On Camina groups of prime power order*, J. Algebra **181** (1996), no. 3, 787–802.

[9] W. Feit and J. G. Thompson, *Solvability of groups of odd order*, Pacific J. Math. **13** (1963), 775–1029.

[10] R. A. Ferraz and J. J. Simón-Pınero, *Central units in metacyclic integral group rings*, Comm. Algebra **36** (2008), no. 10, 3708–3722.

[11] E. G. Goodaire, E. Jespers, and C. Polcino Milies, *Alternative loop rings*, North-Holland Mathematics Studies, vol. 184, North-Holland Publishing Co., Amsterdam, 1996.

[12] G. Higman, *The units of group-rings*, Proc. London Math. Soc. (2) **46** (1940), 231–248.

[13] B. Huppert, *Endliche Gruppen. I*, Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin-New York, 1967.

[14] E. Jespers and Á. del Río, *Group ring groups*, Volume 1: Orders and Generic Constructions of Units, De Gruyter, Berlin-Boston, 2015.

[15] E. Jespers, G. Olteanu, Á. del Río, and I. Van Gelder, *Group rings of finite strongly monomial groups: central units and primitive idempotents*, J. Algebra **387** (2013), 99–116.

[16] ———, *Central units of integral group rings*, Proc. Amer. Math Soc. **142** (2014), 2193–2209.

[17] E. Jespers and M. M. Parmenter, *Construction of central units in integral group rings of finite groups*, Proc. Amer. Math. Soc. **140** (2012), no. 1, 99–107.

[18] E. Jespers, M. M. Parmenter, and S. K. Sehgal, *Central units of integral group rings of nilpotent groups*, Proc. Amer. Math. Soc. **124** (1996), no. 4, 1007–1012.

[19] Y. Li and M. M. Parmenter, *Central units of the integral group ring* $\mathbf{Z}A_5$, Proc. Amer. Math. Soc. **125** (1997), no. 1, 61–65.

[20] Z. Li and J. Hai, *The normalizer property for integral group rings of finite solvable T-groups*, J. Group Theory **15** (2012), no. 2, 237–243.

[21] I. D. Macdonald, *Some p-groups of Frobenius and extra-special type*, Israel J. Math. **40** (1981), no. 3-4, 350–364 (1982).

[22] A. Olivieri, Á. del Río, and J. J. Simón, *On monomial characters and central idempotents of rational group algebras*, Comm. Algebra **32** (2004), no. 4, 1531–1550.

[23] D. S. Passman, *Infinite crossed products*, Pure and Applied Mathematics, vol. 135, Academic Press Inc., Boston, MA, 1989.

[24] I. Reiner, *Maximal orders*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], London-New York, 1975, London Mathematical Society Monographs, No. 5.

[25] J. Ritter and S. K. Sehgal, *Integral group rings with trivial central units*, Proc. Amer. Math. Soc. **108** (1990), no. 2, 327–329.